

Competing structures in ^{188}Pb nucleus

J. Pakarinen,¹ P. Papadakis,² K. Auranen,¹ D. Cox,² T. Grahn,¹ P. Greenlees,¹
Hadynska-Klek,³ A. Herzan,¹ R.-D. Herzberg,² U. Jakobsson,¹ R. Julin,¹ S. Juutinen,¹
J. Konki,¹ M. Leino,¹ A. Mistry,² D. O'Donnell,² P. Peura,¹ P. Ruotsalainen,¹
M. Sandzelius,¹ J. Sarén,¹ J. Sorri,¹ S. Stolze,¹ J. Uusitalo,¹ and K. Wrzosek-Lipska⁴

¹*Department of Physics, University of Jyväskylä,
P.O. Box 35, FI-40014, Jyväskylä, Finland*

²*Department of Physics, Oliver Lodge Laboratory,
University of Liverpool, Liverpool L69 7ZE, United Kingdom*

³*Heavy Ion Laboratory, University of Warsaw,
ul. Pasteura 5A, 02-093 Warszawa, Poland*

⁴*Instituut voor Kern- en Stralingsfysica,
Katholieke Universiteit Leuven, B-3001 Leuven, Belgium*

In the atomic nucleus, the interplay between single-particle motion, collectivity and pairing is seen as a rich tapestry of coexisting nuclear shapes and exotic excitations. One of the richest regions is formed by very neutron-deficient nuclei with the proton number Z close to the magic 82 and the neutron number N close to 104 midshell [1–3]. A considerable body of both theoretical and experimental evidence has been gathered for coexisting configurations possessing different shapes in this region, but yet there are many open questions left. In this presentation, our recent experimental results on ^{188}Pb will be discussed. These include simultaneous in-beam γ -ray and conversion electron spectroscopy using the SAGE spectrometer [4] and Coulomb excitation experiment of post-accelerated ^{188}Pb beam at REX-ISOLDE, CERN [5].

-
- [1] J.L. Wood *et al.*, Phys. Rep. **215**, 101 (1992).
[2] K. Heyde *et al.*, Phys. Rev. C **44**, 2216 (1991).
[3] R. Julin *et al.*, J. Phys. G **27**, R109 (2001).
[4] P. Papadakis *et al.*, AIP Conf. Proc., **1090** (2009) 14.
[5] O. Kester *et al.*, Nucl. Inst. Meth. Phys. Res., Sect. B **204**, 20 (2003).