

## Shape coexistence in mass 40 region studied via E0 and gamma transitions

E. Ideguchi<sup>1</sup>, T. Kibédi<sup>2</sup>, J.T.H. Dowie<sup>2</sup>, T.H. Hoang<sup>1</sup>, M. Kumar Raju<sup>1</sup>, A.A. Akber<sup>2</sup>,  
L. Bignell<sup>2</sup>, B. Coombes<sup>2</sup>, T.K. Eriksen<sup>2</sup>, M.S.M Gerathy<sup>2</sup>, T.J. Gray<sup>2</sup>, G.J. Lane<sup>2</sup>,  
B.P. McCormick<sup>2</sup>, A.J. Mitchell<sup>2</sup>, A.E. Stuchbery<sup>2</sup>, N. Shimizu<sup>3</sup>, Y. Utsuno<sup>4</sup>

<sup>1</sup> RCNP, Osaka University, Ibaraki, Osaka 567-0047, Japan

<sup>2</sup> Department of Nuclear Physics, The Australian National University, ACT 2601, Australia

<sup>3</sup> CNS, The University of Tokyo, Hongo, Bunkyo-ku, Tokyo 113-0033, Japan

<sup>4</sup> ASRC, Japan Atomic Energy Agency, Tokai, Ibaraki 319-1195, Japan

The advent of shape coexistence is a unique feature of atomic nucleus. This phenomenon particularly occurs near spherical closed shell nucleus, where the onset of shape coexistence is based on the balance between stabilizing effect of closed shells to retain spherical shape and the residual interaction which drives the nucleus to deformed shape [1].

The spherical doubly magic nucleus, <sup>40</sup>Ca, is a best example exhibiting such shape coexistence. A unique feature of <sup>40</sup>Ca is an appearance of low-lying 0<sup>+</sup> states. First excited state is 0<sup>+</sup> at 3.3 MeV and the second excited 0<sup>+</sup> state closely locates at 5.2 MeV. These states are understood as band heads of the normal deformed and the superdeformed bands, respectively [2], which corresponds to the multiple shape coexistence in <sup>40</sup>Ca. Similarly, low-lying 0<sup>+</sup> SD band heads are also observed in neighboring nuclei of mass 40 region [3,4,5].

Existence of the superdeformed (SD) band starting from the 0<sup>+</sup> band head is another unique feature of <sup>40</sup>Ca. Although the existence of superdeformed nuclei are reported in many nuclei of various mass regions, A=60, 80, 130, 150, 190 [3], the superdeformed band head 0<sup>+</sup> states are only observed in mass 40 region [4,5], and in the fission isomer region [3]. Such situation makes it difficult to understand the property of superdeformed state, such as the mixing of the states with different configurations. Therefore, <sup>40</sup>Ca is a quite unique nucleus where one can study the electric monopole (E0) transition strength between the band head of superdeformed state and the spherical ground state, which directly reflects the shape mixing [6].

In order to study the property of superdeformed state of <sup>40</sup>Ca, we have performed an experiment to measure the E0 transition from the excited 0<sup>+</sup> states. Experiment was carried out using a <sup>40</sup>Ca(p,p') reaction at the 14UD tandem accelerator facility in Australian National University. The Super-e pair spectrometer [7,8,9], a superconducting magnetic-lens spectrometer, is employed to measure conversion electrons and electron-positron pairs with excellent background suppression. A single germanium detector was also used to measure gamma transitions from the excited states simultaneously.

In the presentation, the experimental results on E0 transition strength from the normal deformed and superdeformed band in <sup>40</sup>Ca and the theoretical studies based on the large-scale shell model calculation will be discussed. Recent results studied via gamma transitions in mass 40 region will be also presented and discussed.

This work is partially supported by the International Joint Research Promotion Program of Osaka University and JSPS KAKENHI Grant Number JP 17H02893.

- [1] K. Heyde and J.L. Wood, Rev. of Mod. Phys. **83**, 1467 (2011)
- [2] E. Ideguchi *et al.*, Phys. Rev. Lett. **87**, 222501 (2001)
- [3] B. Singh, R. Zywina, R.B. Firestone, Nucl. Data Sheets **97**, 241 (2002)
- [4] C.E. Svensson *et al.*, Phys. Rev. Lett. **85**, 2693 (2000)
- [5] E. Ideguchi *et al.*, Phys. Lett. B **686**, 18 (2010)
- [6] J.L. Wood *et al.*, Nucl. Phys. A **651**, 323 (1999)
- [7] T. Kibédi *et al.*, The Astrophysical Journal **489**, 951 (1997)
- [8] L.J. Evitts *et al.*, Phys. Lett. B **779**, 396 (2018)
- [9] L.J. Evitts *et al.*, Phys. Rev. C (in press)